



Molecular insights into hypericin: Unraveling its theoretical potential in Alzheimer's disease intervention

Ivan Vito Ferrari¹ 

¹Institute of Clinical Physiology, National Research Council, Via Aurelia Sud, 54100 Massa, Italy. E-mail: ivanvitoferrari@gmail.com

Article Info

Volume 1, Issue 1, January 2024

Received : 27 October 2023

Accepted : 5 January 2024

Published : 05 January 2024

doi: [10.62587/AFRJMS.1.1.2024.22-24](https://doi.org/10.62587/AFRJMS.1.1.2024.22-24)

Abstract

BACE-1 is involved in the production of beta-amyloid peptides, and inhibiting its activity is a potential therapeutic strategy for managing Alzheimer's disease, as excessive beta-amyloid accumulation is a characteristic feature of the disease. Computational results suggest that hypericin has the potential to act as an inhibitor of BACE-1. The Molecular Docking results indicate a high binding affinity between hypericin and BACE-1, with a notable binding energy score of -10.3 kcal/mol. Additionally, the formation of various chemical bonds, including hydrogen bonds and hydrophobic bonds, further supports the idea that hypericin may effectively interact with the active site of BACE-1. It is important to note that while the *in silico* Molecular Docking results are promising, further *in vitro* and *in vivo* studies would be necessary to validate and better understand the inhibitory potential of hypericin against BACE-1. Additionally, the practical application of hypericin as a BACE-1 inhibitor in a therapeutic context would require careful consideration of factors such as bioavailability, safety, and efficacy.

Keywords: BACE-1, Hypericin, Alzheimer's disease, Molecular docking

© 2024 Ivan Vito Ferrari. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

1. Introduction

Beta-secretase 1 (BACE-1), also known as beta-site amyloid precursor protein cleavage enzyme 1, is an enzyme encoded by the BACE-1 gene in humans. This enzyme is mainly expressed in neurons. Its importance is evident in the formation of myelin sheaths in peripheral nerve cells, with high expression during postnatal stages in mice, related to myelination (Venugopal *et al.*, 2008; Shen *et al.*, 2018; Cervellati *et al.*, 2020; Chacón-Quintero *et al.*, 2021).

The structure of BACE-1 is characterized by two aspartate residues in the active site of its extracellular protein domain. It can function as a dimer, and its cytoplasmic tail is crucial for proper maturation and efficient intracellular trafficking, although it does not directly influence enzymatic activity. It is produced as a proenzyme, undergoing endoproteolysis after leaving the endoplasmic reticulum and reaching the Golgi

* Corresponding author: Ivan Vito Ferrari, Institute of Clinical Physiology, National Research Council, Via Aurelia Sud, 54100 Massa, Italy. E-mail: ivanvitoferrari@gmail.com

apparatus. During this process, the pro-peptide receives additional sugars to increase its molecular mass, and the tail becomes palmitoylated (Venugopal *et al.*, 2008; Shen *et al.*, 2018; Cervellati *et al.*, 2020; Chacón-Quintero *et al.*, 2021).

BACE-1 expression is subject to the influence of the inflammatory state. During Alzheimer's disease (AD), cytokines downregulate PPAR1, an inhibitor of BACE-1 mRNA, thus contributing to the regulation of BACE-1 expression during inflammatory conditions (Venugopal *et al.*, 2008; Shen *et al.*, 2018; Cervellati *et al.*, 2020; Chacón-Quintero *et al.*, 2021). The present study aims to investigate this protein with a natural molecule called Hypericin known for its anti-inflammatory (Dellafiora *et al.*, 2018), anti-viral and anti-tumor properties (Miskovsky, 2002). Several pharmaceutical companies are studying different BACE-1 inhibitors (Moussa-Pacha *et al.*, 2020). However, the priority need is to discover a natural molecule with low side effects that has the ability to behave as a selective inhibitor of this protein, implicated in Alzheimer's disease, a progressive and irreversible neurodegenerative disease that affects the brain. It characterized by a gradual loss of memory and cognitive function, Alzheimer's is the most common form of dementia (Scheltens *et al.*, 2021). The present study is based on computational methods based on Molecular Docking (Fan *et al.*, 2019), a highly precise in-silico technique that aims to automatically calculate the energetic affinities between Hypericin and BACE-1. Furthermore, the amino acids involved in the active site of the protein involved with Hypericin are also investigated.

2. Material and methods

Crystal Structure of Beta-secretase 1 (BACE-1) was taken from Protein Data Bank and Molecular Docking investigation (Fan *et al.*, 2019) was performed by Mcule Database (Odhar *et al.*, 2019). Binding site center of BACE-1 protein (PDB Code 3cic) was X(19,4438), Y(32,672) Z(57,4953). Hypericin was taken from Drug Bank Database.

3. Results and discussion

This brief communication presents, for the first time, an exploration of the interaction between Hypericin and BACE-1. Hypericin, derived from anthraquinone, is naturally found in the yellow flowers of *Hypericum perforatum* (St. John's wort). It demonstrates antidepressant properties, as well as potential antiviral, antineoplastic, and immunostimulating activities (Dellafiora *et al.*, 2018; Miskovsky, 2002).

The selected approach was Molecular Docking (Fan *et al.*, 2019), a well-established and potent method commonly employed in drug design, typically employed prior to conducting *in vitro* and *in vivo* studies.




Figure 1: Displays the docking outcomes of BACE-1 in conjunction with Hypericin within the Ligand Binding Site, as analyzed by Autodock Vina through the Mcule Database. On the left side, 2D diagrams illustrate the residue interactions between the protein and Hypericin. Meanwhile, the right side exhibits the Ligand Binding Site of the protein, highlighting the specific location of Hypericin

In Figure 1, the outcomes of Molecular Docking and associated characterizations between BACE-1 and hypericin are depicted. The results are notably intriguing, revealing a remarkably high binding energy affinity of hypericin with the protein, registering a score of -10.3 kcal/mol. This suggests that, theoretically, this natural substance can effectively bind to the active site of the protein. Furthermore, hypericin seems to establish several chemical bonds with the protein, including hydrogen bonds with ASN 237, THR 236, and ARG 239, as well as hydrophobic bonds with ASP 232, THR 76, TYR 75, and PHE 112.

4. Conclusion

BACE-1's role in beta-amyloid peptide production makes it a target for Alzheimer's disease treatment, aiming to mitigate the associated beta-amyloid accumulation. Computational analyses suggest hypericin's potential as a BACE-1 inhibitor. Molecular Docking reveals a robust binding affinity with a significant energy score of -10.3 kcal/mol, supported by the formation of chemical bonds (hydrogen and hydrophobic) indicating effective interaction at BACE-1's active site. While promising, *in silico* results require validation through *in vitro* and *in vivo* studies to deepen our understanding of hypericin's inhibitory capabilities against BACE-1. Additionally, practical use as a BACE-1 inhibitor mandates careful consideration of factors such as bioavailability, safety, and efficacy in a therapeutic context. Hypericin's capacity to modulate BACE-1 presents a potential avenue for Alzheimer's disease intervention, though further research is vital to confirm its efficacy and safety in more complex biological settings.

References

- Cervellati, C., Trentini, A., Rosta, V., Passaro, A., Bosi, C., Sanz, J. M. and Zuliani, G. (2020). Serum beta-secretase 1 (BACE-1) activity as candidate biomarker for late-onset Alzheimer's disease. *GeroScience*, 42(1); 159-167.
- Chacón-Quintero, M.V., Pineda-López, L.G., Villegas-Lanau, C.A., Posada-Duque, R. and Cardona-Gómez, G. P. (2021). Beta-secretase 1 underlies reactive astrocytes and endothelial disruption in neurodegeneration. *Frontiers in Cellular Neuroscience*, 15; 656832.
- Dellafiora, L., Galaverna, G., Cruciani, G., Dall'Asta, C. and Bruni, R. (2018). On the mechanism of action of anti-inflammatory activity of hypericin: An *in silico* study pointing to the relevance of Janus kinases inhibition. *Molecules*, 23(12); 3058.
- Fan, J., Fu, A. and Zhang, L. (2019). Progress in molecular docking. *Quantitative Biology*, 7; 83-89.
- Miskovsky, P. (2002). Hypericin-a new antiviral and antitumor photosensitizer: mechanism of action and interaction with biological macromolecules. *Current Drug Targets*, 3(1); 55-84.
- Moussa Pacha, N.M., Abdin, S.M., Omar, H.A., Alniss, H. and Al Tel, T.H. (2020). BACE-1 inhibitors: Current status and future directions in treating Alzheimer's disease. *Medicinal Research Reviews*, 40(1), 339-384.
- Odhar, H.A., Rayshan, A.M., Ahjel, S.W., Hashim, A.A. and Albeer, A.A.M.A. (2019). Molecular docking enabled updated screening of the matrix protein VP40 from Ebola virus with millions of compounds in the MCULE database for potential inhibitors. *Bioinformation*, 15(9); 627.
- Scheltens, P., De Strooper, B., Kivipelto, M., Holstege, H., Chételat, G., Teunissen, C.E. and van der Flier, W.M. (2021). Alzheimer's disease. *The Lancet*, 397(10284); 1577-1590.
- Shen, Y., Wang, H., Sun, Q., Yao, H., Keegan, A.P., Mullan, M. and Hampel, H. (2018). Increased plasma beta-secretase 1 may predict conversion to Alzheimer's disease dementia in individuals with mild cognitive impairment. *Biological Psychiatry*, 83(5); 447-455.
- Venugopal, C., Demos, C.M., Jagannatha Rao, K.S., Pappolla, M.A. and Sambamurti, K. (2008). Beta-secretase: structure, function, and evolution. *CNS & Neurological Disorders-Drug Targets (Formerly Current Drug Targets-CNS & Neurological Disorders)*, 7(3); 278-294.

Cite this article as: Ivan Vito Ferrari. (2024). Molecular insights into hypericin: Unraveling its theoretical potential in Alzheimer's disease intervention. *African Research Journal of Medical Sciences*. 1(1), 22-24. doi: 10.62587/AFRJMS.1.1.2024.22-24.